



[Kapoor, et al.](#)

Impact of PHGG on constipation prevention: A systematic review and meta-analysis. *Journal of Functional Foods*, 33 (2017) 52-66.

[Carlson, J. Slavin, J., et al.](#)

In vitro analysis of PHGG fermentation on identified gut microbiota. *Anaerobe*. 42, pp60-66, 2016

[Carlson, J. Slavin, J., et al.](#)

In vitro analysis of PHGG fermentation differences between six individuals. *Food Funct*, 7, pp1833, 2016

[Hung T, et al.](#)

Dietary Fermentable Fiber Reduces Intestinal Barrier Defects and Inflammation in Colitic Mice. *J Nutr*. 146, pp1970-1979, 2016

[Horii, Y, et al.](#)

PHGG enhances colonic epithelial wound healing via activation of RhoA and ERK1/2. *Food & Function*. 13, pp3176-83, 2016.

[Takagi, T., et al.](#)

PHGG ameliorates intestinal inflammation. *British journal of Nutrition*. 116, pp1199-1205, 2016

[Rao, T.P](#)

Role of guar fiber in appetite control. *Physiology & Behavior*. 164, pp277-283, 2016

[Kapoor, et al.](#)

Soluble dietary fibre partially hydrolyzed guar gum markedly impacts on postprandial hyperglycaemia, hyperlipidaemia and incretins metabolic hormones over time in healthy and glucose intolerant subjects. *Journal of Functional Foods* 24, pp207-220, 2016

[Tokunaga, M, et al.](#)

Effect of PHGG on Postprandial hyperglycemia – A randomized double-blind, placebo controlled crossover study. *Jpn Pharmacol Ther*. 44, pp185-91, 2016.

[Niv, E, et al.](#)

Randomized clinical study: PHGG versus placebo in the treatment of patients with IBS. *Nutrition & Metabolism*. 13, pp10, 2016

[Alam NH, et al.](#)

Efficacy of partially hydrolyzed guar gum (PHGG) supplemented modified oral rehydration solution in the treatment of severely malnourished children with watery diarrhoea: a randomised double-blind controlled trial. *J Health Popul Nutr*. 34, pp3, 2016

[Naito, Y., et al.](#)

Identification of cysteinylated transthyretin, a predictive biomarker of treatment response to partially hydrolyzed guar gum in type 2 diabetes rats, by surface-enhanced laser desorption/ionization time-of-flight mass spectrometry. *J Clin Biochem Nutr*. 58, pp23-33, 2016

[Russo L, et al.](#)

PHGG in the treatment of IBS with constipation” Effects of gender, age, and BMI. *Gastroenterology*. 21, pp104-10, 2015

[Rao T.P., et al.](#)

Post-meal perceivable satiety and subsequent energy intake with intake of partially hydrolyzed guar gum. *British Journal of Nutrition*. 113, pp1489-98, 2015.

[Ohashi, Y., et al.](#)

Consumption of partially hydrolysed guar gum stimulates Bifidobacteria and butyrate-producing bacteria in the human large intestine. *Beneficial Microbes*. 6, pp451-455, 2015

[Brillantino, A, et al.](#)

Maintenance Therapy with Partially Hydrolyzed Guar Gum in the Conservative Treatment of Chronic Anal Fissure: Results of a Prospective, Randomized Study. *BioMed Research International*. 2014.

[Sevinc AI, et al.](#)

Improvement of colonic healing by preoperative oral partially hydrolyzed guar gum in rats which underwent preoperative radiotherapy. *Journal of Drug Targeting*. 22, pp262-268, 2014



[Ishihara N, et al.](#)

Physiological effect and applications of partially hydrolyzed guar gum. *Luminacoids* 17, pp11-18, 2013

[Paul SP, et al.](#)

Challenges in Management of Irritable Bowel Syndrome in Children. *Ind Pediatr* 50, pp1137-1143, 2013

[Salaria Ammeta, et al.](#)

Prebiotics in the microencapsulating matrix enhance the viability of probiotic *Lactobacillus acidophilus* LA1. *Int J Fermented Foods*. 2, pp33-45, 2013

[Dall'Alba, V., et al.](#)

Improvement of the metabolic syndrome profile by soluble fibre – guar gum – in patients with type 2 diabetes: a randomized clinical trial. *British Journal of Nutrition*. 110, pp1601–1610, 2013

[Quartarone, Q., et al.](#)

Role of PHGG as a dietary fiber: a review article. *Minerva Gastroenterologica E Dietologica*. 59, pp329 -340, 2013.

[Romano C, et al.](#)

Partially hydrolyzed guar gum in pediatric functional abdominal pain. *World Journal of Gastroenterology*. 19, pp235-240, 2013

[Finley J, et al.](#)

Safety Assessment and Caloric Value of Partially Hydrolyzed Guar Gum. *J. Agric Food Chem* 61, pp1756-1771, 2013

[Ohashi Y, et al.](#)

Faecal fermentation of partially hydrolyzed guar gum, *J. Funct. Foods*. 4, pp398-402, 2012

[Yasukawa, et al.](#)

Partially hydrolyzed guar gum affects the expression of genes involved in host defense functions and cholesterol absorption in colonic mucosa of db/db male mice, *J. Clin. Biochem. Nutr.* 51, pp33–38, 2012

[Furnari M, et al.](#)

Efficacy of partially hydrolyzed guar gum in reducing methane excretion and clinical manifestation of subjects suffering from irritable bowel syndrome. *Gastroenterology Abs Sa2058*. 2012

[Ishihara N, et al.](#)

Normalization of evacuation on the saline in a nursing home for elderly by partially hydrolyzed guar gum. *Journal for the integrated study of dietary habits*. 22, pp315-319, 2012

[Maeda H, et al.](#)

Partially hydrolyzed guar gum intake ameliorates constipation, improves nutritional status and reduces indoxylsulfuric acid in dialysis patients. *Kidney Research and Clinical Practice* 31, A53. 2012

[Prasad PS, et al.](#)

Stool Consistency and Abdominal Pain in Irritable Bowel Syndrome May Be Improved by Partially Hydrolysed Guar Gum. *Journal of Pediatric Gastroenterology & Nutrition*, 53, pp582-583, 2011

[Furnari M, et al.](#)

Clinical trial: the combination of rifaximin with partially hydrolysed guar gum is more effective than rifaximin alone in eradicating small intestinal bacterial overgrowth, *Aliment. Pharmacol. Ther.* 32, pp1000–1006, 2010

[Ustunda G, et al.](#)

Can partially hydrolyzed guar gum be an alternative to lactulose in treatment of childhood constipation?, *Turk. J. Gastroenterol.* 21, pp360-364, 2010

[Kuo DC, et al.](#)

Partially hydrolyzed guar gum supplement reduces high-fat diet increased blood lipids and oxidative stress and ameliorates FeCl₃-induced acute arterial injury in hamsters, *J. Biomed. Sci.* 2, 16, 15, 2009

[Takahashi T, et al.](#)

Hydrolyzed guar gum decreases postprandial blood glucose and glucose absorption in the rat small intestine, *Nutr Res.* 29, pp419-425, 2009



[Lluch A, et al.](#)

Short-term appetite-reducing effects of a low-fat dairy product enriched with PHGG. *Food Quality and Preference* 21. *Food Quality and Preference* 21, pp402-409, 2009

[Atila K, et al.](#)

Partially hydrolyzed guar gum attenuates the severity of pouchitis in a rat model of ileal J pouch-anal anastomosis, *Dig. Dis. Sci.* 54, pp522-529, 2009

[Yoon S-J, et al.](#)

Chemical and physical properties, safety and application of partially hydrolyzed guar gum as dietary fiber, *J. Clin. Biochem. Nutr.* 42, pp1-7, 2008

[Alam NH, et al.](#)

Efficacy of partially hydrolyzed guar gum-added oral rehydration solution in the treatment of severe cholera in adults, *Digestion.* 78, pp24-29, 2008

[Nakamura, S, et al.](#)

Suppressive effect of partially hydrolyzed guar gum on transitory diarrhea induced by ingestion of maltitol and lactitol in healthy humans. *European Journal of Clinical Nutrition.* pp1-8, 2007.

[Maenaka, T, et al.](#)

Effects of partially hydrolyzed guar gum on postprandial blood glucose level and disaccharidase, *J.JSMUFF*, 4, 3, pp195-201, 2007

[Giannini, E, et al.](#)

Role of partially hydrolyzed guar gum in the treatment of irritable bowel syndrome, *Nutrition.* 22, pp334-342, 2006

[Stewart, M, et al.](#)

Molecular weight of guar gum affect SCFA profile in model intestinal fermentation, *Mol. Nutr. Food Res.* 50, pp971-976, 2006.

[Naito, Y, et al.](#)

Prevention of life style related disease by dietary fiber, *J. JSMUFF.* 5, pp271-275, 2006.

[Naito, Y, et al.](#)

Partially hydrolyzed guar gum down-regulates colonic inflammatory response in dextran sulfate sodium-induced colitis in mice, *J. Nutr. Biochem.* 17, pp402-409, 2006.

[Cassia Freitas, K, et al.](#)

Partially hydrolyzed guar gum increases intestinal absorption of iron in growing rats with iron deficiency anemia, *Clin. Nutr.* 25, pp851-858, 2006.

[Yoon, SJ, et al.](#)

Physiological functions of partially hydrolyzed guar gum, *J Clin Biochem Nutr.* 39, pp134-144, 2006.

[Alam, N, et al.](#)

Partially hydrolysed guar gum supplemented comminuted chicken diet in persistent diarrhoea. A randomised controlled trial, *Arch. Dis. Child.* 90, pp195-199, 2005.

[Parisi, G, et al.](#)

Treatment effects of partially hydrolyzed guar gum on symptoms and quality of life of patients with irritable bowel syndrome. A multicenter randomized open trial, *Dig. Dis. Sci.* 50, pp1107-1112, 2005.

[Minekus, M, et al.](#)

Effect of partially hydrolyzed guar gum (PHGG) on the bioaccessibility of fat and cholesterol, *Biosci. Biotechnol. Biochem.* 69, pp932-938, 2005.

[Ishihara, N](#)

Sunfiber[®], a water soluble dietary fiber, improves intestinal microflora balance, *FOOD Style* 21 9, pp53-56, 2005. (Japanese)



[Pklkas, AM, et al.](#)

Comparison of different fibers for in vitro production of short chain fatty acids by intestinal microflora, *J. Med. Food.* 8, pp113-116, 2005.

[Yokozawa, T, et al.](#)

Green tea polyphenols and dietary fibre protect against kidney damage in rats with diabetic nephropathy, *J. Pharm. Pharmacol.* 57, pp773-780, 2005.

[Rushidi, TA, et al.](#)

Control of diarrhea by fiber-enriched diet in ICU patients on enteral nutrition: A prospective randomized controlled trial, *Clin. Nutr.* 23, pp1344-1352, 2004.

[Cihan, A, et al.](#)

Comparison of early postoperative enteral nutrients versus chow on colonic anastomotic healing in normal animals, *Eur Surg Res.* 36, pp112-115, 2004.

[Bosaeus, I, et al.](#)

Fibre effects of intestinal functions (diarrhoea, constipation and irritable bowel syndrome), *Clin. Nutr. Suppl.* 1, pp33-389, 2004.

[Suzuki, T, et al.](#)

Ingestion of guar gum hydrolysate, a soluble and fermentable nondigestible saccharide, improves glucose intolerance and prevents hypertriglyceridemia in rats fed fructose, *J. Nutr.* 134, pp1942-1947, 2004.

[Trinidad, T, et al.](#)

Glycemic index of Sunfiber (*Cyamopsis tetragonolobus*) products in normal and diabetic subjects, *Int. J. Food Sci. Technol.* 39, pp1093-1098, 2004.

[Bar, A, et al.](#)

Reducing the glycemic impact of food. A new role for some dietary fibres, *Innovations in Food Technol.*, Feb, pp34-38, 2004.

[Suzuki, T, et al.](#)

Ingestion of guar gum hydrolysate, a soluble and fermentable nondigestible saccharide, improves glucose intolerance and prevents hypertriglyceridemia in rats fed fructose, *J. Nutr.* 134, pp1942-1947, 2004.

[Nakao, M, et al.](#)

Usefulness of soluble dietary fiber for the treatment of diarrhea during enteral nutrition in elderly patients, *JSPEN*, 18, pp53-57, 2003.

[Slavin, JL, et al.](#)

Partially hydrolyzed guar gum: clinical nutrition uses, *Nutrition*, 19, pp549-552, 2003.

[YeunHwa, G, et al.](#)

Effect of enzyme hydrolyzed guar gum on elevation of blood glucose levels after meal, *Medecine ad biology.* 47, pp19-24, 2003.

[Ishihara, N](#)

Improvement effect of partially hydrolyzed guar gum, *Food Style*21, 7, pp79-82, 2003.

[Yamada, K, et al.](#)

Effect of dietary fiber on the lipid metabolism and immune function of aged Sprague-Dawley rats, *Biosci. Biotechnol. Biochem.* 67, pp429-433, 2003.

[Nakao, M, et al.](#)

Usefulness of soluble dietary fiber for the treatment of diarrhea during enteral nutrition in elderly patients, *Nutrition.* 18, pp35-39, 2002.

[Parisi, G, et al.](#)

High-fiber diet supplementation in patients with irritable bowel syndrome (IBS): A multicenter, randomized, open trial comparison between wheat bran diet and partially hydrolyzed guar gum (PHGG), *Dig. Dis. Sci.* 47, pp1697-1704, 2002.

[Nakao, M, et al.](#)

Usefulness of soluble dietary fiber for the treatment of diarrhea during enteral nutrition in elderly patients. *Nutrition.* 18, pp35-39, 2002.



[Giaccari, S, et al.](#)

Partially hydrolyzed guar gum: a fiber as coadjuvant in the irritable colon syndrome. *Clin Ter.* 152, pp21-5, 2001

[Spapen, H, et al.](#)

Soluble fiber reduces the incidence of diarrhea in septic patients receiving total enteral nutrition: a prospective, double-blind, randomized, and controlled trial. *Clinical Nutrition.* 20, pp301-305, 2001.

[Tuohy, K.M., et al.](#)

The prebiotic effects of biscuits containing partially hydrolyzed guar gum and fructo-oligosaccharides – a human volunteer study. *British Journal of Nutrition.* 86, pp341-348, 2001.

[Alam, N, et al.](#)

Partially hydrolyzed guar gum-supplemented oral rehydration solution in the treatment of acute diarrhea in children. *Journal of Pediatric Gastroenterology and Nutrition.* 31, pp503-507, 2000.

[Ishihara, N, et al.](#)

Preventative effect of partially hydrolyzed guar gum on infection of salmonella enteritidis in young and laying hens. *Poultry Science* 79, pp689-697, 2000.

[Greenberg, N.A.](#)

The chemical, physical, and physiological properties of partially hydrolyzed guar gum. Novartis Nutr. Corp. AACC 2000 annual meeting. Nov 5-9, 2000.

[Tanaka, T, et al.](#)

Effects of rice gruel containing partially hydrolyzed guar gum on human defecation. *Journal of Nutritional Food.* 3, pp45-52, 2000.

[Velazquez, M, et al.](#)

Effect of Oligosaccharides and Fibre substitutes on short chain fatty acid production by human faecal microflora. *Anaerobe.* 6, pp87-92, 2000.

[Sarianom C, et al.](#)

Long-term fiber intervention program: Reduction in enema use at a developmental care facility. *Supplement* 100, ppA-82, 2000.

[Yomada, k, et al.](#)

Dietary effect of guar gum and its partially hydrolyzed product on the lipid metabolism and immune function of Sprague-dawley rats. *Biosci. Biotechnol. Biochem.* 63, pp2163-2167, 1999.

[Chu, et al.](#)

Chemical and functional properties of partially hydrolyzed guar gum (Sunfiber) as a dietary Fibre. *Innovations in Food Technology.* Feb, pp9-14, 1999.

[Okazaki, H, et al.](#)

Improvement in defecation by a beverage containing partially hydrolyzed guar gum. *Journal of Nutritional Food,* 2, pp1-8, 1999.

[Hara, H, et al.](#)

Ingestion of guar-gum hydrolysate partially restores calcium absorption in the large intestine lowered by suppression of gastric acid secretion in rats. *British Journal of Nutrition.* 81, pp315-321, 1999.

[Hara, H, et al.](#)

Ingestion of guar gum hydrolysate, a soluble fiber, increases calcium absorption in totally gastrectomized rats. *American Society for Nutritional Sciences. J Nutr.* 107, pp1340-1348, 1999.

[Alam, N.H., et al.](#)

Effects of a partially hydrolyzed guar gum on intestinal absorption of carbohydrate, protein, and fat: a double-blind controlled study in volunteers. *Clinical Nutrition* 17, pp125-129, 1998.

[Patrick, P, et al.](#)

Effect of supplements of partially hydrolyzed guar gum on the occurrence of constipation and use of laxative agents. *Journal of the American dietetic association.* 98, pp912-914, 1998.



[Greenberg, N.A., et al.](#)

Partially hydrolyzed guar gum as a source of fiber. *Cereal Foods World*. 43, pp703-707, 1998.

[Heini, A.F., et al.](#)

Effect of hydrolyzed guar fiber on fasting and postprandial satiety and satiety hormones: a double-blind, placebo-controlled trial during controlled weight loss. *International Journal of Obesity*. 22, pp906-909, 1998.

[Tsuda, K, et al.](#)

Effect of partially hydrolyzed guar gum on elevation of blood glucose after sugar intake in human volunteers. *Journal of Japanese Association for dietary fiber research*. 2, pp15-22, 1998.

[Yamatoya, K, et al.](#)

Effects of hydrolyzed guar gum on cholesterol and glucose in humans. *Food Hydrocolloids*. 11, pp239-242, 1997.

[Favier, M, et al.](#)

The cholesterol-lowering effect of guar gum is not the result of a simple diversion of bile acids toward fecal excretion. *Lipids*. 32, pp953-959, 1997.

[Peters, A.](#)

Addition of hydrolyzed guar to enteral feeding products in type I diabetic patients. *Diabetes Care*. 19, pp899-900, 1996.

[Weaver, G, et al.](#)

Dietary guar gum alters colonic Microbial fermentation in azoxymethane-treated rats. *American Institute of Nutrition*, pp1979-1991, 1996.

[Hara, H, et al.](#)

Increases in calcium absorption with ingestion of soluble dietary fibre, guar gum hydrolysate, depend on the caecum in partially nephrectomized and normal rats. *British Journal of Nutrition*. 76, pp773-784, 1996.

[Furuse, M, et al](#)

Effects of partially hydrolyzed guar gum on feeding behavior and crop emptying rate in chicks. *British Poultry Science*. 37, pp223-227, 1996.

[Takahashi, H, et al.](#)

Effect of liquid diets with or without partially hydrolyzed guar gum on intestinal microbial flora and function of rats. *Nutritional Research*. 15, pp527-536, 1995.

[Yamatoya, K, et al.](#)

Effect of hydrolyzed guar gum on frequency and feeling of defecation in humans. *Oyo Toshitsu Kagaku*. 42, pp251-257, 1995.

[Golay, A, et al.](#)

The effect of a liquid supplement containing guar gum and fructose on glucose tolerance in non-insulin-dependent diabetic patients. *Nutr. Melab Cardiovasc Dis*. 5, pp141-148, 1995.

[Takahashi, H, et al.](#)

Influence of partially hydrolyzed guar gum on constipation in women. *J. Nutr. Sci. Vitaminol.*, 40, pp251-259, 1994.

[Takahashi, H, et al](#)

Influence of intact and partially hydrolyzed guar gum on iron utilization in rats fed on iron-deficient diets. *Comp. Biochem. Physiol.* 109, pp75-82, 1994.

[Okubo, T, et al.](#)

Effects of partially hydrolyzed guar gum intake on human intestinal microflora and its metabolism. *Biosci. Biotech. Biochem.* 58, pp1364-1369, 1994.

[Homman, H, et al.](#)

Reduction in diarrhea incidence by soluble fiber in patients receiving total or supplemental enteral nutrition. *Journal of Parenteral and External Nutrition*. 18, pp486-490, 1994.



[Vandeven, M, et al.](#)

Effects of liquid preloads with difference fructose/fibre concentrations on subsequent food intake and ratings or hunger in women. *Appetite*, 23, pp139-146, 1994.

[Takahashi, H, et al.](#)

Protein and energy utilization of growing rats fed on the diets containing intact or partially hydrolyzed guar gum. *Comp. Biochem. Physiol.* 107A, pp255-260, 1993.

[Yamatoyo, K, et al.](#)

Effects of partially hydrolyzed guar gum on postprandial plasma glucose and lipid levels in humans. *Nippon Eiyo Syokuryo Gakkaishi*. 46, pp199-203, 1993.

[Meier, R, et al.](#)

Effect of a liquid diet with and without soluble fiber supplementation on intestinal transit and cholecystokinin release in volunteers. *Journal of Parenteral and Enteral Nutrition*. 17, pp231-236, 1993.

[Takahashi, H, et al.](#)

Effect of partially hydrolyzed guar gum on fecal output in human volunteers. *Nutritional Research*, 13, pp649-657, 1993.

[Lampe, J, et al.](#)

Gastrointestinal effects of modified guar gum and soy polysaccharide as part of an enteral formula diet. *Journal of Parenteral and Enteral Nutrition*. 16, pp538-546, 1992.

[Ide, T, et al.](#)

Hypolipidemic effects of guar gum and its enzyme hydrolysate in rats fed highly saturated fat diets. *Ann Nutr Metab*. 35, pp34-44, 1991.

[Takeno, F, et al.](#)

Effect of partially decomposed guar gum on high-cholesterol-fed rats and non-dietary fiber-fed rats. *Research Laboratories, Dainippon Pharmaceutical Co., Ltd*, 43, pp421-425, 1990.